

MRS Fall 1996 Meeting
December 2-6, 1996
Boston MA

Submitted to Symposium FF: Dynamics in Small Confining Systems IV

Binding of Confined Vortex-Pairs in Micron-Scale, Patterned Co Films*

Peter J. Bedrossian and Stephen P. Vernon
Lawrence Livermore National Laboratory
Livermore CA 94551

ABSTRACT

Using Magnetic Force Microscopy (MFM) and Vibrating Sample Magnetometry (VSM), we have characterized the magnetic microstructures of micron-scale, Co ellipses which were patterned from Co films with thicknesses up to 70 nm. A variety of sizes and shapes (2x8, 2x4, 4x8, and 4x16 micron ellipses, and 2 and 4-micron diameter circles) show no remanant magnetization after saturation along their short axes. MFM shows that the absence of remanance results from closure domain patterns arising from the presence of vortices. Each elliptical size and shape is associated with a specific number and pattern of vortices which appear at remanance. The presence of the boundary results in demagnetization effects, which strongly favor tangential orientation of domains at the ellipse edges.

In 2x8 and 4x8 micron ellipses, two vortices with opposite circulations appear on the major axis at remanance, with their separation determined by a minimum in the potential well defined by vortex-vortex and vortex-edge interactions. Application of a magnetic field along the hard axis results in antisymmetric displacement of the vortices along the easy axis, and the vortex-vortex separation scales linearly with the applied magnetic field.

*Research was performed under the auspices of the US Department of Energy at Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.